

(19)

Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) **EP 0 818 454 A1**

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
14.01.1998 Bulletin 1998/03

(51) Int Cl.⁶: **C07D 401/04**, C07D 413/14,
A61K 31/445

(21) Application number: 97500099.3

(22) Date of filing: 03.06.1997

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE**
Designated Extension States:
SI

(30) Priority: 04.06.1996 ES 9601236


(71) Applicant: **FABRICA ESPANOLA DE
PRODUCTOS
QUIMICOS Y FARMACEUTICOS, S.A. (FAES)**
E-48940 Lejona-Lamiaco (Vizcaya) (ES)

(72) Inventors:
• **Orjales, Aurelio**
48990-Neguri, Vizcaya (ES)
• **Rubio, Victor**
48990 Getxo, Vizcaya (ES)
• **Bordell, Maravillas**
48940 Leioa, Vizcaya (ES)

(74) Representative: **Flaccus, Rolf-Dieter, Dr.**
Patentanwalt
Sperlingsweg 32
50389 Wesseling (DE)

(54) **Benzimidazole derivatives with antihistaminic activity**

(57) New benzimidazole derivatives of formula:



in which R_1 is H or a short chain hydrocarbon group such as methyl, ethyl, isopropyl, cyclopropyl, vinyl, etc., and R_2 is a group selected from among the following: CH_2OH , $COOH$, $COOR_3$ and 4,4-dimethyl-2-oxazoliny, R_3 being a short chain alkyl group such as methyls ethyl, etc., are described.


A description is also made of the preparation of these compounds, which have a high H_1 antihistaminic and antiallergic activity and are devoid of effects on the central nervous and cardiovascular systems.

Description**Object of the Invention**

5 The present invention relates to new benzimidazole derivatives with H₁ antihistaminic activity, lacking cardiotoxic effects.

Background of the Invention

10 The prior art closest to the compounds of the present invention is in Spanish patent No. 9201512 which describes a number of piperidine benzimidazole derivatives with antihistaminic and antiallergic activity of general formula:



20 The major structural difference between the compounds of the present invention and those of the said patent is the presence of oxygenated functions in the phenyl group substitution. An important pharmacophorous character has moreover been found for these oxygenated functions which consists essentially in a selectivity of action and provides a pharmacological profile distinct from that of other known antihistamines. The compounds disclosed in this invention present an almost exclusive H₁ antihistaminic pharmacological activity and are therefore devoid of action on other pharmacological receptors even at doses much higher than the therapeutic ones. Because of this selectivity in action, they are valuable instruments in treating allergic-type conditions, particularly allowing their unrestricted use by persons under any other concomitant medication whatsoever, and in the case of patients with pathological cardiocirculatory disturbances.

Description of the Invention

35 It has long since been known that histamine plays a very important role in allergic-type diseases, such as allergic rhinitis, conjunctivitis, urticaria and asthma; antihistaminic compounds acting at the H₁-receptor histamine level are useful for treating such conditions.


First generation H₁ antihistamines presented a number of adverse effects, such as sedation and dryness of the mouth, resulting from its action on the central nervous system and colinergic receptors.

40 The search for molecules that would not cross the haematoencephalic barrier brought about the displacement of the early antihistamines by other second generation antihistamines which overcame the side effects linked to their action on the central nervous system. This new generation of antihistamines, amongst which noteworthy, due to their extensive use worldwide, are terfenadine and astemizole, has recently displayed a negative aspect in the form of dangerous cardiovascular effects, extending the QT space and ventricular arrhythmia, which has required its use to be avoided in those cases in which the patient is prone to suffering such disturbances or when he is being treated with substances that may interfere with his metabolism.

45 Attempts at obtaining safe and efficient H₁ antihistamines have multiplied in recent years and this research has resulted in several recent patent applications claiming pharmaceutical compositions for treating allergic diseases containing antihistamines devoid of arrhythmogenic effects, which is the case of US patent application number 924,156 (3/08/92) and international patent application number 95/00480 (05/01/95).

50 The present invention relates to a group of new compounds with benzimidazolic structure having potent selective H₁ antihistaminic activity, lacking activity on the central nervous system and on the cardiovascular system.


The compounds subject of the present invention have the following general formula:




I

in which R_1 is hydrogen or a short chain hydrocarbon group such as methyl, ethyl, isopropyl, cyclopropyl or vinyl, and R_2 is a group selected from among CH_2OH , COOH , COOR_3 and 4,4-dimethyl-2-oxazolanyl, R_3 being a short chain alkyl group as previously defined, and their addition salts with pharmacologically acceptable acids or bases.


Compounds I in which R_1 is a short chain alkyl group and R_2 is the 4,4-dimethyl-2-oxazolanyl group can be conveniently prepared by an N-alkylation reaction of 2-(4-piperidiny)-1H-benzimidazole with an alkylating agent of formula III wherein X is a good leaving group in nucleophilic substitution reactions such as Cl, Br, I, R_4SO_2 , R_2SO_3 , etc., in the presence of an inorganic base, such as an alkaline metal carbonate or bicarbonate within an organic solvent, followed by another N-alkylation reaction of the resulting benzimidazole IV with an ether of formula $\text{XCH}_2\text{CH}_2\text{OR}_1$, wherein X has the meaning given above and R_1 is a short chain hydrocarbon group, such as Me, Et, i-Pr, cyclopropyl, vinyl, etc., in the presence of a hydride or an alkaline metal carbonate.





II



III




Ia





IV

Compounds I, in which R_1 is a short chain alkyl group and R_2 is a carboxyl group, can be conveniently prepared by hydrolysis of the benzimidazoles Ia with a mineral acid such as HCl or H_2SO_4 .




Ia



Ib


Compounds I in which R_1 is a short chain alkyl group and R_2 is a COOR_3 group in which R_3 is a short chain alkyl group,

can be prepared by acid hydrolysis of the benzimidazoles Ia in the presence of an alcohol solvent R_3OH , in adequate conditions for transesterification.




15

Compounds I in which R_1 is a short chain alkyl group and R_2 is a CH_2OH group can be prepared by a reduction of the benzimidazoles Ib or Ic with a suitable reducing agent, such as aluminium and lithium hydride.




30

Compound I in which R_1 is a hydrogen and R_2 is a 4,4-dimethyl-2-oxazoliny group can be prepared by an alkylation of the benzimidazole IN with ethyl chloroacetate in the presence of a hydride or an alkaline metal carbonate to yield the ester V, which is then reduced by a reducing agent such as aluminium and lithium hydride.



55

Compound I in which R_1 is a hydrogen and R_2 is a $COOH$ group is conveniently prepared by hydrolysis with a mineral acid such as HCl or H_2SO_4 of the compound Id.



The obtained new benzimidazoles I can be turned into pharmaceutically acceptable salts by treatment with suitable acids or bases.

Compounds of formula I have useful pharmacological properties. In particular, they are potent H_1 antihistamines. This activity was clearly demonstrated *in vitro* by blocking the histamine-induced contractions in the isolated guinea pig ileum (Magnus, Pflügers, Arch. Ges. Physiol., 102, 123 (1904); Arunlakshana, O. and Schild, H.O., Br. J. Pharmacol. 14, 48-58 (1959)) and *in vivo* by the capacity to inhibit the increment of histamine-induced cutaneous capillary permeability in rats (Lefebvre, P., Salmon, J., Leconte and Cauwenberge, V.H., C.R. Soc. Biol. 156, 183-186 (1962); Udaka, K., Takeuchi, Y. and Morat, H.Z., Proc. Soc. Exp. Biol. Med. 133, 1384-1387 (1970)).

Thus, compound Ib (R_1 =ethyl) proved to be a potent guinea pig ileum H_1 -receptor histamine mixed antagonist, with calculated $pA_2=7.98-8.10$ and $pD_2'=6.50$. This same compound inhibited *in vivo* the increase in capillary permeability in rats with a DE_{50} close to 2 mg/kg p.o. At doses of 5 mg/kg p.o. it maintained a significant activity, in excess of 50%, for at least 6 hours.

These compounds are highly selective in their pharmacological action, and present no significant anticholinergic activity nor activity on the central nervous and cardiovascular systems. Thus, compound Ib (R_1 =Et) is not able to antagonize significantly the acetylcholine-induced contractions in isolated guinea pig ileum at 0.1 M concentrations and does not modify the spontaneous motor activity of the rat at 100 mg/kg p.o.; furthermore, this same compound, administered at 20 mg/kg i.v., induces no morphological ECG disturbance nor does it increase the QT_c interval in rats.

In view of their useful pharmacological antihistaminic and antiallergic properties, the compounds described in the present invention can be formulated in several pharmaceutical forms to be later administered orally, topically, injectably and rectally. Oral preparations are made by intimately mixing a quantity effective as antihistaminic of one of the products described in the present invention with excipients such as lactose, cellulose, talc and the like for tablets or capsules, or water, glycols, alcohols, oils and the like for syrups, solutions and suspensions. Topical administration can be made in the form of creams, ointments, gels, solutions and transcutaneous plasters, using agents such as vaseline, polyethylene glycols, etc. as a carrier. In preparations for injectables, the excipient will be, at least for the most part, sterilised water, although other excipients, such as saline solutions, glucose solutions, etc., or mixtures thereof, may be added to enhance solubility.

The examples detailed below illustrate the present invention without howsoever limiting its scope.

Example 1

Preparation of 1-(2-ethoxyethyl)-2-[1-(2-(4-(1-(4,4-dimethyl- Δ^2 -oxazoline-2-yl)-1-(methylethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole. (Ia, R_1 =Et)

3.57 g of sodium carbonate were added to a suspension of 14 g 2-(4-(1-(4,4-dimethyl- Δ^2 -oxazoline-2-yl)-1-methylethyl)phenyl)ethyl p-toluene-sulphonate and 6.78 g 2-(4-piperidinyl)-1H-benzimidazole in 60 ml of DMF and the resulting suspension was heated at 80° for 14 hours. The DMF was concentrated and the reaction mass was poured onto water/ice whereupon a solid crystallised which was filtered, washed with water and dried at 50°C to yield 10 g of 2-[1-(2-(4-(1-(4,4-dimethyl- Δ^2 -oxazoline-2-yl)-1-methylethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole. The resulting solid was dissolved in 25 ml of DMF and 1.2 g of a sodium hydride in 60% oil suspension was added to this. The resulting suspension was stirred at room temperature for two hours and 2.44 g of 2-chloroethylethylether were added. The reaction mass was heated at 80°C for 16 hours, cooled, poured on water/ice, extracted with ether and washed with water and with saturated sodium chloride solution. The ethereal solution was dried over anhydrous sodium sulphate and concentrated to yield 11.2 g of 1-(2-ethoxyethyl)-2-[1-(2-(4-(1-(4,4-dimethyl- Δ^2 -oxazoline-2-yl)-1-methylethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole.

MP: 98-100°C (ethanol).

RMN- 1H ($CDCl_3$), δ : 1.1 (t, 3H); 1.3 (s, 6H); 1.5 (s, 6H); 1.9 (m, 2H); 2.1 (m, 4H); 2.6 (t, 2H); 2.8 (t, 2H); 3.0 (m, 1H); 3.1 (d, 2H); 3.4 (c, 2H); 3.7 (t, 2H); 3.9 (s, 2H); 4.3 (t, 2H); 7.1-7.3 (m, 7H); 7.7-7.8 (m, 1H).

RMN- ^{13}C ($CDCl_3$), δ : 14.96; 27.38; 28.15; 31.06; 33.10; 34.53; 40.18; 43.60; 53.71; 60.46; 66.74; 66.83; 68.59; 79.14;

109.09; 119.41; 121.71; 121.88; 125.30; 128.73; 134.78; 138.72; 142.72; 143.04; 158.41 and 177.70.

Example 2

Preparation of 2-[4-(2-(4-(1-(2-ethoxyethyl)benzimidazole-2-yl) piperidine-1-yl)ethyl)phenyl]-2-methylpropanoic acid.
(Ib, R₁=Et)

6.72 g of 1-(2-ethoxyethyl)-2-[1-(2-(4-(1-(4,4-dimethyl-Δ²-oxazoline-2-yl)-1-methylethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole (Ia) were dissolved in 170 ml of HCl 3N and refluxed for an hour. This was cooled and taken to pH 7 with 50% sodium hydroxide. The solution was extracted with *n*-butanol, washed with water, dried over anhydrous sodium sulphate and concentrated. Methanol (30 ml) and 50% sodium hydroxide (40 ml) were added to the residue and refluxed for thirty minutes. The methanol was distilled off and water was added until dissolution was complete. This was extracted with ether and the aqueous layer was taken to pH 7 with 20% HCl and saturated with sodium chloride, whereupon a solid precipitated which was filtered, washed repeatedly with water and dried in a vacuum dryer at 50°C to yield 3.5 g of 2-[4-(2-(4-(1-(2-ethoxyethyl) benzimidazole-2-yl)piperidine-1-yl)ethyl)phenyl]-2-methylpropanoic acid.

MP: 199-201°C

RMN-¹H (DMSO-d₆), δ: 1.0 (t, 3H); 1.4 (s, 6H); 1.8 (m, 4H); 2.2 (m, 2H); 2.5 (t, 2H); 2.7 (t, 2H); 3.0 (m, 3H); 3.3 (c, 2H); 3.6 (t, 2H); 4.4 (t, 2H); 7.0-7.3 (m, 6H); 7.4-7.6 (m, 2H).

RMN-¹³C (DMSO-d₆), δ: 14.90; 26.59; 30.97; 32.22; 33.39; 43.04; 45.50; 53.08; 60.05; 65.70; 68.43; 110.18; 118.40; 121.16; 121.35; 125.47; 128.42; 134.72; 138.33; 142.29; 143.03; 158.60; and 177.87.

Example 3

Preparation of ethyl 2-[4-(2-(4-(1-(2-ethoxyethyl)benzimidazole-2-yl) piperidine-1-yl)ethyl)phenyl]-2-methylpropanoate.

(Ic, R₁=ET, R₃=ET) Concentrated sulphuric acid (20 ml) were added over a solution of 10 g of 1-(2-ethoxyethyl)-2-[1-(2-(4-(1-(4,4-dimethyl-Δ²-oxazoline-2-yl)-1-methylethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole in 250 ml of ethanol, and this was refluxed for 16 hours. This was cooled and 1 litre of ether was added. The organic layer was separated and washed with water, 10% sodium bicarbonate solution and once again with water. This was dried over anhydrous sodium sulphate and concentrated to yield 7 g of an oil which was purified by flash-chromatography using a 95/5 chloroform/ethanol mixture as eluent to yield 5 g of ethyl 2-[4-(2-(4-(1-(2-ethoxyethyl) benzimidazole-2-yl)piperidine-1-yl)ethyl)phenyl]-2-methylpropanoate in the form of an oil.

RMN-¹H (CDCl₃), δ: 1.1 (t, 3H); 1.2 (t, 3H); 1.5 (s, 6H); 2.0 (m, 2H); 2.2 (m, 4H); 2.6 (t, 2H); 2.8 (t, 2H); 3.0 (m, 1H); 3.2 (m, 2H); 3.4 (c, 2H); 3.7 (t, 2H); 4.1 (c, 2H); 4.3 (t, 2H); 7.1-7.3 (m, 7H); 7.6-7.7 (m, 1H).

RMN-¹³C (CDCl₃), δ: 13.86; 14.80; 26.35; 30.62; 32.73; 33.87; 43.48; 45.91; 53.26; 60.11; 60.49; 66.61; 68.40; 109.02; 119.16; 121.55; 121.75; 125.40; 128.50; 134.56; 138.40; 142.29; 142.51; 158.13 and 176.53.

Example 4

Preparation of 1-(2-ethoxyethyl)-2-[1-(2-(4-(1,1-dimethyl-2-hydroxyethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole. (Id, R₁=Et)

1 g of aluminium and lithium hydride was dissolved in 30 ml of THF and 3 g of ethyl 2-[4-(2-(4-(1-(2-ethoxyethyl) benzimidazole-2-yl) piperidine-1-yl)ethyl)phenyl]-2-methylpropanoate were added dropwise thereto. This was stirred for four hours at room temperature and some millilitres of water were added to eliminate excess hydride. The solution was filtered and the filtrate was washed with a saturated sodium chloride solution. This was dried and concentrated. The residue was redissolved in chloroform and washed with water, dried and concentrated. The residue was purified by flash-chromatography using a hexane/ether/isopropyl-amine mixture (2/7.5/0.5) as an eluent, to yield 1.5 g of 1-(2-ethoxyethyl)-2-[1-(2-(4-(1,1-dimethyl-2-hydroxyethyl) phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole.

MP: 112-114°C

RMN-¹H (CDCl₃), δ: 1.0 (t, 3H); 1.4 (s, 6H); 1.9-2.1 (m, 2H); 2.1-2.3 (m, 4H); 2.6 (t, 2H); 2.8 (t, 2H); 3.0 (m, 1H); 3.2 (d, 2H); 3.4 (c, 2H); 3.6 (s, 2H); 3.7 (t, 2H); 4.3 (t, 2H); 7.1-7.4 (m, 7H); 7.8 (m, 1H). RMN-¹³C (CDCl₃), δ: 15.01; 25.34; 31.07; 33.07; 34.53; 39.78; 43.64; 53.72; 60.52; 66.88; 68.62; 73.07; 109.13; 119.44; 121.77; 121.94; 126.22; 128.80; 134.78; 138.39; 142.71; 143.90; 158.45.

Example 5Preparation of 1-(2-hydroxyethyl)-2-[1-(2-(4-(1-(4,4-dimethyl- Δ^2 -oxazoline-2-yl)-1-methylethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole. (Ie)

5g of 2-[1-(2-(4-(1-(4,4-dimethyl- Δ^2 -oxazoline-2-yl)-1-methylethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole were dissolved in 30 ml of DMF and 0.54 g of a sodium hydride in oil suspension were added thereto. The resulting suspension was stirred for two hours at room temperature and 1.19 ml of ethyl chloroacetate were added dropwise. The reaction mass was heated at 70° for 16 hours, cooled and poured on 300 ml of water. This was extracted with ether and the ethereal layer was washed with water, dried over anhydrous sodium sulphate and filtered. 0.8 g of aluminium and lithium hydride were dissolved in 30 ml of ether and the previously filtered ethereal phase was added dropwise to this solution. This was stirred for 4 hours at room temperature, and 20 ml of a 10% sodium hydroxide solution were added thereto. This was saturated with sodium chloride and the ethereal layer was separated. The aqueous phase was extracted with ether. The ethereal phases were all blended together and washed with water and with a saturated sodium chloride solution. This was dried over anhydrous sodium sulphate and concentrated to yield 2.6 g of 1-(2-hydroxyethyl)-2-[1-(2-(4-(1-(4,4-dimethyl- Δ^2 -oxazoline-2-yl)-1-methylethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole in the form of an oil.

RMN-¹H (CDCl₃), δ : 1.3 (s, 6H); 1.6 (s, 6H); 1.8-2.2 (m, 6H); 2.6 (t, 2H); 2.8 (t, 2H); 2.9 (m, 1H); 3.0-3.1 (m, 2H); 3.7 (s, 2H); 4.0 (s, 2H); 4.3 (t, 2H); 7.1-7.4 (m, 7H); 7.7 (m, 1H).

RMN-¹³C (CDCl₃), δ : 15.01; 25.34; 31.07; 33.07; 34.53; 39.78; 43.64; 53.72; 60.52; 66.88; 68.62; 73.07; 109.13; 119.44; 121.77; 121.94; 126.22; 128.80; 134.78; 138.39; 142.71; 143.90; 158.45.

Example 6Preparation of 2-[4-(2-(4-(1-(2-hydroxyethyl)benzimidazole-2-yl)piperidine-1-yl)ethyl)phenyl]-2-methylpropanoic acid. (II)


5 g of 1-(2-hydroxyethyl)-2-[1-(2-(4-(1-(4,4-dimethyl- Δ^2 -oxazoline-2-yl)-1-methylethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole (Ie) were dissolved in 45 ml of 3N HCl and refluxed for an hour. This was taken to a basic pH with 50% NaOH and 20 ml of ethylene glycol were added. This was heated at 190°C for three hours with simultaneous distillation and then concentrated in vacuo. Water was added and extracted with ether. The aqueous layer was taken to pH 7 with diluted HCl, saturated with sodium chloride and extracted with *n*-butanol. The ethereal extract was dried and concentrated. The residue was recrystallised in acetone/methanol to yield 2.7 g of 2-[4-(2-(4-(1-(2-hydroxyethyl)benzimidazole-2-yl)piperidine-1-yl)ethyl)phenyl]-2-methylpropanoic acid.

MP: 218°C (breaks down)

RMN-¹H (CDCl₃), δ : 1.4 (s, 6H); 2.0-2.1 (m, 4H); 2.7-2.9 (m, 4H); 2.9-3.1 (t, 2H); 3.2-3.5 (m, 3H); 3.7 (t, 2H); 4.3 (t, 2H); 6.9-7.1 (m, 2H); 7.1-7.2 (m, 2H); 7.2-7.3 (m, 2H); 7.3-7.4 (m, 1H); 7.4-7.5 (m, 1H).

Claims

1. New benzimidazole derivatives of formula:



in which R₁ is hydrogen or a short chain hydrocarbon group such as methyl, ethyl, isopropyl, cyclopropyl or vinyl, and R₂ is a group selected from among CH₂OH, COOH, COOR₃ and 4,4-dimethyl-2-oxazolinyl, R₃ being a short chain alkyl group as defined hereinabove, and their addition salts with pharmacologically acceptable acids or bases.

2. A compound in accordance with claim 1 which is 1-(2-ethoxy ethyl)-2-[1-(2-(4-(1-(4,4-dimethyl- Δ^2 -oxazoline-2-yl)-1-methylethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole or an addition salt with a pharmaceutically acceptable acid or base.

EP 0 818 454 A1

3. A compound in accordance with claim 1 which is 2-[4-(2-(4-(1-(2-ethoxyethyl)benzimidazole-2-yl)piperidine-1-yl)ethyl)phenyl]-2-methyl propanoic acid or an addition salt with a pharmaceutically acceptable acid or base.
- 5 4. A compound in accordance with claim 1 which is ethyl 2-[4-(2-(4-(1-(2-ethoxyethyl)benzimidazole-2-yl)piperidine-1-yl)ethyl)phenyl]-2-methylpropanoate or an addition salt with a pharmaceutically, acceptable acid or base.
5. A compound in accordance with claim 1 which is 1-(2-ethoxyethyl)-2-[1-(2-(4-(1,1-dimethyl-2-hydroxy-ethyl)phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole or an addition salt with a pharmaceutically acceptable acid or base.
- 10 6. A compound in accordance with claim 1 which is 1-(2-hydroxyethyl)-2-[1-(2-(4-(1-(4,4-dimethyl- Δ^2 -oxazoline-2-yl)-1-methylethyl) phenyl)ethyl)piperidine-4-yl]-1H-benzimidazole or an addition salt with a pharmaceutically acceptable acid or base.
- 15 7. A compound in accordance with claim 1 which is 2-[4-(2-(4-(1-(2-hydroxyethyl)benzimidazole-2-yl)piperidine-1-yl)ethyl)phenyl]-2-methyl propanoic acid or an addition salt with a pharmaceutically acceptable acid or base.
8. An antihistaminic pharmaceutical composition, characterised by containing an effective quantity as antihistaminic of one of the compounds claimed in claims 1-7 as an active ingredient, mixed with one or several excipients.
- 20 9. A process for treating allergic diseases in patients, which comprises administering a pharmaceutical composition in accordance with claim 8, at suitable doses.

25

30

35

40

45

50

55



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 97 50 0099 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Y	EP 0 580 541 A (FABRICA ESPANOLA DE PRODUCTOS QUIMICOS Y FARMACEUTICOS, S.A.) * claims 1,7 *	1,8	C07D401/04 C07D413/14 A61K31/445
D	& ES 2 048 109 A ---		
Y	WO 94 03170 A (SEPRACOR INC., ET AL.) * abstract; claims *	1,8	
D	& US appl. ser. no. 924156 (03.08.92) ---		
D,Y	WO 95 00480 A (MERRELL DOW PHARMACEUTICALS INC.) * claim 1 *	1	
A	JOURNAL OF HETEROCYCLIC CHEMISTRY, vol. 24, 1987, pages 31-7, XP002041046 R. IEMURA ET AL.: "Synthesis of Benzimidazole Derivatives as Potential H1-Antihistaminic Agents" ---	1,8	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07D
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search		Date of completion of the search	Examiner
BERLIN		18 September 1997	Hass, C
CATEGORY OF CITED DOCUMENTS			
<p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>I : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EP 1 FORM 150 (1.1.91) (P0403)



European Patent Office

EP 97 50 0099 - C -

INCOMPLETE SEARCH

The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.

Claims searched completely:
Claims searched incompletely:
Claims not searched:

Reason for the limitation of the search: Although claim 9 is directed to a method of treatment of (diagnostic method practised on) the human/animal body (Article 52(4) EPC) the search has been carried out and based on the alleged effects of the compound/composition.

European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 97 50 0099

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	JOURNAL OF HETEROCYCLIC CHEMISTRY, vol. 32, 1995, pages 707-18, XP002041047 A. ORJALES ET AL.: "Synthesis and Structure-Activity Relationship of New Piperidinyl and Piperazinyl Derivatives as Antiallergics" * page 707 * -----	1,8	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)

EPO FORM 1503 01.01 (P04C10)